

Neutrino Mass Matrix with Two Zeros and Leptogenesis

Satoru Kaneko ^{*} and Morimitsu Tanimoto [†]

Department of Physics, Niigata University, Ikarashi 2-8050, 950-2181 Niigata, JAPAN

ABSTRACT

The leptogenesis is studied in the neutrino mass matrix with two zeros, which reduces the number of independent phases of the CP violation. These texture zeros are decomposed into the Dirac neutrino mass matrix and the right-handed Majorana one in the see-saw mechanism. It is found that there are six textures for the Dirac neutrino mass matrix, among which one texture with three zeros has three phases, three textures with four zeros have two phases, and two textures with five zeros have only one phase in the basis with the real right-handed Majorana mass matrix. The relation between the leptogenesis and the low energy CP violation is discussed in typical textures. It is remarked that the leptogenesis is linked strongly with the CP violation in the neutrino oscillations. The predicted baryon asymmetry of the universe is consistent with the observed value.

^{*}E-mail address: kaneko@muse.sc.niigata-u.ac.jp

[†]E-mail address: tanimoto@muse.sc.niigata-u.ac.jp

The Super-Kamiokande has almost confirmed the neutrino oscillation in the atmospheric neutrinos, which favors the $\nu_\mu \rightarrow \nu_\tau$ process [1]. For the solar neutrinos [2, 3], the recent data of the Super-Kamiokande and the Sudbury Neutrino Observatory also favor strongly the neutrino oscillation $\nu_e \rightarrow \nu_{\mu,\tau}$ with the large mixing angle (LMA) MSW solution [4, 5]. These results indicate the neutrino masses and mixings [6], especially, the bi-large flavor mixing. It is therefore possible to investigate the textures of the neutrino mass matrix.

Recently, Frampton, Glashow and Marfatia [7] found seven acceptable textures of the neutrino mass matrix with two independent vanishing entries in the basis with the diagonal charged lepton mass matrix. The further studies of these textures were presented by some authors [8, 9, 10, 11]. Another insight has been also given focusing on texture zeros [12]. Since these textures are given for the light left-handed neutrinos, one needs to find the see-saw realization [13] of these textures from the standpoint of the model building. We have found the Dirac neutrino mass matrices and the right-handed Majorana neutrino ones, which give those seven textures [11].

These textures should be tested in the future experiments. We have already shown that they can be tested in the lepton flavor violating processes, especially, $\mu \rightarrow e\gamma$ [11]. In order to search another test of them, we discuss the leptogenesis in this paper. We study the relation between the leptogenesis and the low energy CP violation in these textures. We also predict the magnitude of the baryon asymmetry of the universe.

The leptogenesis [14] is an interesting mechanism to explain the observed baryon asymmetry of the universe. Since the leptogenesis depends strongly on the texture and the CP violating phases of the neutrino mass matrix, it is expected that the leptogenesis links with the low energy neutrino phenomena, especially the low energy CP violation in the neutrino oscillations [15, 16, 17]. The CP violation required for the leptogenesis stems from the CP violating phases in the right-handed sector, whereas the CP violation in the neutrino oscillations [18] can be described by the left-handed neutrino mixing matrix. In general, the independent CP violating phases are six for three generations and without a left-handed Majorana mass term, and therefore it is very difficult to find links between the leptogenesis and the low energy CP violation. Reduction of the number of phases has been done by considering the two right-handed

neutrinos [19, 20]. The interesting connection between the leptogenesis and the low energy CP violation has been presented in those works. It is remarked that there is one massless neutrino in this scheme.

In order to reduce the number of phases, we propose another approach, which is the texture zeros with the three right-handed neutrinos, being originated from the work by Frampton, Glashow and Marfatia [7]. We show that the low energy leptonic CP violation can be affected by the CP violating phases responsible for the leptogenesis in these texture zeros.

Let us discuss the structure of the CP violating phases in the neutrino mass matrix. There are seven acceptable textures with two independent zeros for the effective neutrino mass matrix M_ν [7]. The two textures (A_1 and A_2) correspond to the hierarchical neutrino mass spectrum while others ($B_1 \sim B_4$ and C) correspond to the quasi-degenerate neutrino mass spectrum. In this paper, we concentrate only the hierarchical case of the textures A_1 and A_2 :

$$A_1 : M_\nu = \begin{pmatrix} \mathbf{0} & \mathbf{0} & \times \\ \mathbf{0} & \times & \times \\ \times & \times & \times \end{pmatrix}, \quad A_2 : M_\nu = \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \times & \times \\ \mathbf{0} & \times & \times \end{pmatrix}, \quad (1)$$

where the charged lepton mass matrix is the diagonal one. Since the component $M_\nu(1,1)$ is zero in these textures, the amplitude for the neutrinoless double beta decay vanishes to lowest order in neutrino masses.

In principle these textures are given at the low energy scale because the experimental data are put to determine zeros, however, the structure of zeros in the mass matrix is not changed by the one-loop renormalization group equations [21]. Therefore, we discuss the see-saw realization in these textures at the right-handed Majorana neutrino mass scale:

$$\mathbf{M}_\nu = \mathbf{m}_\mathbf{D} \mathbf{M}_\mathbf{R}^{-1} \mathbf{m}_\mathbf{D}^T, \quad (2)$$

where $\mathbf{m}_\mathbf{D}$ and $\mathbf{M}_\mathbf{R}$ are the Dirac neutrino mass matrix and the right-handed Majorana neutrino one, respectively. As far as we exclude the possibility that zeros are originated from cancellations among coefficients in the see-saw mechanism, the see-saw realization of these seven textures are not trivial. Then, these zeros should come from zeros of the Dirac neutrino mass matrix and the right-handed Majorana neutrino one. These results are summarized in the reference [11].

Once the right-handed Majorana neutrino mass matrix is specified, the Dirac neutrino ones

are selected in order to reproduce the textures in eq.(1). The simplest right-handed Majorana neutrino mass matrix is the diagonal one, but there is no solution to give desirable texture zeros unless we consider cancellations between matrix elements of the Dirac and Majorana ones. Therefore, we take a following texture for the right-handed Majorana neutrinos

$$\mathbf{M}_R = \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \times & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \times \end{pmatrix}, \quad (3)$$

where $\times$'s denote non-zero entries. It should be noted that specifying the texture of the right-handed Majorana neutrinos is a choice of weak basis. Furthermore we can take the matrix in eq.(3) to be real. Then, the CP violating phases appear only in the Dirac neutrino mass matrix. In this case, we have six Dirac neutrino mass matrices, which reproduce the texture A_1 in eq.(1):

$$\mathbf{m}_D = \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \mathbf{0} & \times & \times \\ \times & \times & \times \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \mathbf{0} & \times & \times \\ \times & \mathbf{0} & \times \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \mathbf{0} & \times & \times \\ \times & \times & \mathbf{0} \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \mathbf{0} & \times & \times \\ \times & \mathbf{0} & \mathbf{0} \end{pmatrix}, \quad (4)$$

and

$$\mathbf{m}_D = \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \times \\ \times & \times & \times \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \times \\ \times & \mathbf{0} & \times \end{pmatrix}, \quad (5)$$

where $\times$'s denote complex numbers. As seen later the textures in eq.(5) give different predictions from the ones in eq.(4) for the decay rate of $\mu \rightarrow e\gamma$.

For the texture A_2 in eq.(1), we obtain

$$\mathbf{m}_D = \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \times & \times \\ \mathbf{0} & \times & \times \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \times & \times \\ \mathbf{0} & \mathbf{0} & \times \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \times & \mathbf{0} \\ \mathbf{0} & \times & \times \end{pmatrix}, \quad (6)$$

and

$$\mathbf{m}_D = \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \mathbf{0} & \times \\ \mathbf{0} & \times & \times \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \mathbf{0} & \times \\ \mathbf{0} & \mathbf{0} & \times \end{pmatrix}, \quad \begin{pmatrix} \mathbf{0} & \times & \mathbf{0} \\ \times & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \times & \times \end{pmatrix}. \quad (7)$$

The textures in eq.(7) also give different predictions from the ones in eq.(6) for $\mu \rightarrow e\gamma$.

Although there are phases in all non-zero entries, those phases are not necessarily physical quantities. Actually, three phases are removed by the re-definition of the left-handed neutrino fields. There is no freedom of the re-definition for the right-handed ones in the basis with real $\mathbf{M}_R$. As seen in eqs.(4), (5), (6), (7), the number of independent phases are classified as: one

texture of three zeros with three phases, three textures of four zeros with two phases and two textures of five zeros with one phase. Thus, we get the reduction of the number of phases in our textures while there are six independent phases in the neutrino mass matrix without zeros.

What kind of experiments can test these textures? It is well known that the Yukawa coupling of the neutrino contributes to the lepton flavor violation (LFV). Many authors have studied the LFV in the minimal supersymmetric standard model (MSSM) with right-handed neutrinos assuming the relevant neutrino mass matrix [22]~[27]. In the MSSM with soft breaking terms, there exist lepton flavor violating terms such as off-diagonal elements of slepton mass matrices and trilinear couplings (A-term). It is noticed that large neutrino Yukawa couplings and large lepton mixings generate the large LFV in the left-handed slepton masses. For example, the decay rate of $\mu \rightarrow e\gamma$ can be approximated as follows:

$$\Gamma(\mu \rightarrow e\gamma) \simeq \frac{e^2}{16\pi} m_\mu^5 F \left| \frac{(6 + 2a_0^2)m_{S0}^2}{16\pi^2} (\mathbf{Y}_\nu \mathbf{Y}_\nu^\dagger)_{21} \ln \frac{M_X}{M_R} \right|^2, \quad (8)$$

where the neutrino Yukawa coupling matrix $\mathbf{Y}_\nu$ is given as $\mathbf{Y}_\nu = \mathbf{m}_\mathbf{D}/v_2$ (v_2 is a VEV of Higgs) at the right-handed mass scale M_R , and F is a function of masses and mixings for SUSY particles. In eq.(8), we assume the universal scalar mass (m_{S0}) for all scalars and the universal A-term ($A_f = a_0 m_{S0} Y_f$) at the GUT scale M_X . Therefore the branching ratio $\mu \rightarrow e\gamma$ depends considerably on the texture of the Dirac neutrino [25, 26, 27].

The magnitude of $(\mathbf{m}_\mathbf{D} \mathbf{m}_\mathbf{D}^\dagger)_{21}$ is important to predict the branching ratio of $\mu \rightarrow e\gamma$ ¹. Many works have shown that this branching ratio is too large [25, 26]. However, zeros in the Dirac neutrino mass matrix may lead to $(\mathbf{m}_\mathbf{D} \mathbf{m}_\mathbf{D}^\dagger)_{21} = 0$, and then it suppress the $\mu \rightarrow e\gamma$ decay. The branching ratio is safely predicted to be below the present experimental upper bound 1.2×10^{-11} [28] due to zeros. Actually, the case of $(\mathbf{m}_\mathbf{D} \mathbf{m}_\mathbf{D}^\dagger)_{21} = 0$ was examined carefully in ref. [29]. We have several textures, which lead to $(\mathbf{m}_\mathbf{D} \mathbf{m}_\mathbf{D}^\dagger)_{21} = 0$ as seen in eqs.(5) and (7). Therefore we examine the textures of the Dirac neutrinos in eqs.(5) and (7) for the leptogenesis in the following.

At first, let us discuss the texture in the A_1 case in eq.(1). Key ingredients are the structure

¹Although the right-handed Majorana neutrino mass matrix is not a diagonal one in our case, $\mathbf{m}_\mathbf{D} \mathbf{m}_\mathbf{D}^\dagger$ does not depend on the right-handed sector.

of CP violating phases and the magnitudes of each entry for the Dirac neutrino mass matrices in eq.(5) since the right-handed Majorana neutrino mass matrix is taken to be real. Although the non-zero entries $\times$ in the Dirac neutrino mass matrix are complex, three phases are removed by the re-definition of the left-handed neutrino fields. There is no freedom of re-definition for the right-handed ones in the basis with real $\mathbf{M}_R$. Furthermore, we move to the diagonal basis of the right-handed Majorana neutrino mass matrix in order to calculate the magnitude of the leptogenesis. Then, the Dirac neutrino mass matrices $\overline{\mathbf{m}}_D$ in the new basis is given as follows:

$$\overline{\mathbf{m}}_D = \mathbf{P}_L \mathbf{m}_D \mathbf{O}_R , \quad (9)$$

where $\mathbf{P}_L$ is a diagonal phase matrix and $\mathbf{O}_R$ is the orthogonal matrix which diagonalizes $\mathbf{M}_R$ as $\mathbf{O}_R^T \mathbf{M}_R \mathbf{O}_R$ in eq.(3). Since the phase matrix $\mathbf{P}_L$ can remove one phase in each row of $\mathbf{m}_D$, three phases disappear in $\overline{\mathbf{m}}_D$.

By taking three eigenvalues of $\mathbf{M}_R$ as follows ²:

$$M_1 = -\lambda^m M_0 , \quad M_2 = \lambda^n M_0 , \quad M_3 = M_0 , \quad (10)$$

where $\lambda \simeq 0.22$, we obtain the orthogonal matrix $\mathbf{O}_R$ as

$$\mathbf{O}_R = \begin{pmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix} , \quad \tan^2 \theta = \lambda^{m-n} . \quad (11)$$

We take $m = 4$, $n = 2$ as a typical case in order to find links between the leptogenesis and the low energy CP violation. The magnitude of each entry in M_ν (A_1) is given to be consistent with the experimental data as follows:

$$M_\nu = m_0 \begin{pmatrix} 0 & 0 & \lambda \\ 0 & 1 & 1 \\ \lambda & 1 & 1 \end{pmatrix} , \quad (12)$$

where $m_0 \simeq \sqrt{\Delta m_{\text{atm}}^2}/2$ with $\Delta m_{\text{atm}}^2 \simeq 2.5 \times 10^{-3} \text{eV}^2$ is taken. Then two mass matrices $\mathbf{m}_D$ in eq.(5) can be parametrized in the new basis as follows:

$$\overline{\mathbf{m}}_D = m_{D0} \begin{pmatrix} 0 & a\lambda^2 & 0 \\ 0 & 0 & b \\ c\lambda^2 e^{i\rho} & d\lambda^k e^{i\sigma} & f \end{pmatrix} O_R, \quad (k \geq 1) \quad (\text{A11}); \quad m_{D0} \begin{pmatrix} 0 & a\lambda^2 & 0 \\ 0 & 0 & b \\ c\lambda^2 e^{i\rho} & 0 & f \end{pmatrix} O_R, \quad (\text{A12}) \quad (13)$$

²The minus sign of M_1 is necessary to reproduce $\mathbf{M}_R$ in eq.(3). This minus sign is transferred to $\mathbf{m}_D$ by the right-handed diagonal phase matrix $\text{diag}(i, 1, 1)$.

where k is a positive integer with $k \geq 1$, and a, b, c, d, f are real and order one coefficients. The magnitude of m_{D0} is determined by the relation $m_{D0}^2 \simeq m_0 M_0$. There are two independent phases in (A11), but only one phase in (A12) as seen in eq.(13). The texture in (A12) corresponds to the $d = 0$ limit in (A11).

In the limit $M_1 \ll M_2, M_3$, the lepton number asymmetry (CP asymmetry) for the lightest heavy Majorana neutrino (N_1) decays into $l^\mp \phi^\pm$ [30] is given by

$$\epsilon_1 = \frac{\Gamma_1 - \overline{\Gamma}_1}{\Gamma_1 + \overline{\Gamma}_1} \simeq -\frac{3}{16\pi v^2} \left(\frac{\text{Im}[\{(\overline{m}_D^\dagger \overline{m}_D)_{12}\}^2] M_1}{(\overline{m}_D^\dagger \overline{m}_D)_{11} M_2} + \frac{\text{Im}[\{(\overline{m}_D^\dagger \overline{m}_D)_{13}\}^2] M_1}{(\overline{m}_D^\dagger \overline{m}_D)_{11} M_3} \right), \quad (14)$$

where $v = 174$ GeV. The lepton asymmetry Y_L is related to the CP asymmetry through the relation

$$Y_L = \frac{n_L - n_{\overline{L}}}{s} = \kappa \frac{\epsilon_1}{g_*}, \quad (15)$$

where s denotes the entropy density, g_* is the effective number of relativistic degrees of freedom contributing to the entropy and κ is the so-called dilution factor which accounts for the washout processes (inverse decay and lepton number violating scattering). In the case of the standard model (SM), one gets $g_* = 106.75$.

The produced lepton asymmetry Y_L is converted into a net baryon asymmetry Y_B through the $(B + L)$ -violating sphaleron processes. One finds the relation [31]

$$Y_B = \xi Y_{B-L} = \frac{\xi}{\xi - 1} Y_L, \quad \xi = \frac{8 N_f + 4 N_H}{22 N_f + 13 N_H}, \quad (16)$$

where N_f and N_H are the number of fermion families and complex Higgs doublets, respectively. Taking into account $N_f = 3$ and $N_H = 1$ in the SM, we get

$$Y_B \simeq -\frac{28}{51} Y_L. \quad (17)$$

On the other hand, the low energy CP violation, which is a measurable quantity in the long baseline neutrino oscillations [18], is given by the Jarlskog determinant J [32], which is calculated by

$$\det[\mathbf{M}_\ell \mathbf{M}_\ell^\dagger, \mathbf{M}_\nu \mathbf{M}_\nu^\dagger] = -2iJ(m_\tau^2 - m_\mu^2)(m_\mu^2 - m_e^2)(m_e^2 - m_\tau^2)(m_3^2 - m_2^2)(m_2^2 - m_1^2)(m_1^2 - m_3^2), \quad (18)$$

where $\mathbf{M}_\ell$ is the diagonal charged lepton mass matrix, and m_1, m_2, m_3 are neutrino masses.

It is very interesting to investigate links between the leptogenesis (ϵ_1) and the low energy CP violation (J) in the textures of eq.(13). In the texture (A11), ϵ_1 and J are given as

$$\begin{aligned} \epsilon_1 &\simeq -\frac{3m_{D0}^2}{16\pi v^2} \left[d^2 \lambda^{2(k+1)} \frac{c^2 \sin 2(\rho - \sigma) - 2cd\lambda^{k-1} \sin(\rho - \sigma)}{c^2 + d^2 \lambda^{2(k-1)} - 2cd\lambda^{k-1} \cos(\rho - \sigma)} \right. \\ &\quad \left. + f^2 \lambda^4 \frac{c^2 \sin 2\rho + d^2 \lambda^{2(k-1)} \sin 2\sigma - 2cd\lambda^{k-1} \sin(\rho + \sigma)}{c^2 + d^2 \lambda^{2(k-1)} - 2cd\lambda^{k-1} \cos(\rho - \sigma)} \right], \\ &\hspace{25em} (k \geq 1) \end{aligned} \quad (19)$$

$$J \simeq \frac{1}{64} a^2 b^4 c^3 f^2 \lambda^2 [c \sin 2\rho - 2d\lambda^{k-1} \sin(\rho + \sigma)] \frac{\Delta m_{\text{atm}}^2}{\Delta m_{\text{sol}}^2},$$

respectively. Both ϵ_1 and J are given in terms of two phases ρ and σ . However, the dominant terms are given by phase ρ as far as $k \geq 2$. It is remarked that the leptogenesis is linked strongly with the CP violation in the neutrino oscillations. The relative sign of ϵ_1 and J is opposite in this case. The case of $k = 1$ is exceptional one. Detailed numerical study will be needed in $k = 1$ because both ϵ_1 and J depend on parameters c , d , ρ and σ .

The case of the texture (A12) is clear because there is only one phase ρ . In this case, ϵ_1 and J are given in the $d = 0$ limit of the texture (A11) as follows:

$$\epsilon_1 \simeq -\frac{3m_{D0}^2}{16\pi v^2} f^2 \lambda^4 \sin 2\rho, \quad (20)$$

$$J \simeq \frac{1}{64} a^2 b^4 c^4 f^2 \lambda^2 \frac{\Delta m_{\text{atm}}^2}{\Delta m_{\text{sol}}^2} \sin 2\rho.$$

The relative sign of ϵ_1 and J is also opposite. It may be helpful to note that only the second term in the right-handed side of eq.(14) contributes to ϵ_1 since $\text{Im}[\{(\overline{m}_D^\dagger \overline{m}_D)_{12}\}^2]$ vanishes.

On the other hand, taking the LMA-MSW solution $\Delta m_{\text{sol}}^2 / \Delta m_{\text{atm}}^2 = \lambda^3 \sim \lambda^2$ and $\sin 2\rho \simeq 1$, we predict $J \simeq 0.01$, which is rather large and then is favored for the experimental measurement.

Next, consider the case A₂ for M_ν . Since the magnitude of each entry in M_ν is given to be consistent with the experimental data as follows:

$$M_\nu = m_0 \begin{pmatrix} 0 & \lambda & 0 \\ \lambda & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}, \quad (21)$$

three mass matrices $\mathbf{m}_D$ in eq.(7) can be parametrized in the new basis as follows:

$$\overline{\mathbf{m}}_D = m_{D0} \begin{pmatrix} 0 & a\lambda^2 & 0 \\ b\lambda^2 & 0 & ce^{i\rho} \\ 0 & d\lambda^p e^{i\sigma} & f \end{pmatrix} O_R, \quad (\text{A21}); \quad m_{D0} \begin{pmatrix} 0 & a\lambda^2 & 0 \\ b\lambda^2 & 0 & c\lambda^q e^{i\rho} \\ 0 & d\lambda e^{i\sigma} & f \end{pmatrix} O_R, \quad (\text{A22}) \quad (22)$$

where $p \geq 1$ and $q \geq 0$, and

$$\overline{\mathbf{m}}_{\mathbf{D}} = m_{D0} \begin{pmatrix} 0 & a\lambda^2 & 0 \\ b\lambda^2 & 0 & ce^{i\rho} \\ 0 & 0 & f \end{pmatrix} O_R, \quad (\text{A23}); \quad m_{D0} \begin{pmatrix} 0 & a\lambda^2 & 0 \\ b\lambda^2 & 0 & 0 \\ 0 & d\lambda e^{i\sigma} & f \end{pmatrix} O_R, \quad (\text{A24}). \quad (23)$$

The textures in (A23) and (A24) correspond to the $d = 0$ and $c = 0$ limits of (A21) and (A22) in eq.(22), respectively.

For the texture (A21) ϵ_1 and J are given as

$$\epsilon_1 \simeq \frac{3m_{D0}^2}{16\pi v^2} \frac{1}{b^2 + d^2 \lambda^{2(p-1)}} \lambda^4 [b^2 c^2 \sin 2\rho - d^2 f^2 \lambda^{2(p-1)} \sin 2\sigma - 2bcd f \lambda^{p-1} \sin(\rho - \sigma)], \quad (p \geq 1) \quad (24)$$

$$J \simeq \frac{1}{64} a^2 b^3 f^2 \lambda^2 \frac{\Delta m_{\text{atm}}^2}{\Delta m_{\text{sol}}^2} [bc^2 f^2 \sin 2\rho + d\lambda^{p-1} \{2c^3 f \sin(\rho - \sigma) + bc^2 d \lambda^{p-1} \sin 2(\rho - \sigma) + b^3 d \lambda^{p-1} \sin 2\sigma + 2b^2 c f \sin(\rho + \sigma)\}],$$

and for the texture (A22) ϵ_1 and J are given as

$$\epsilon_1 \simeq \frac{3m_{D0}^2}{16\pi v^2} \frac{1}{b^2 + d^2} \lambda^4 [b^2 c^2 \lambda^{2q} \sin 2\rho - d^2 f^2 \sin 2\sigma - 2bcd f \lambda^q \sin(\rho - \sigma)], \quad (q \geq 0) \quad (25)$$

$$J \simeq \frac{1}{64} a^2 b^3 f^2 \lambda^2 \frac{\Delta m_{\text{atm}}^2}{\Delta m_{\text{sol}}^2} [b^3 d^2 \sin 2\sigma + c\lambda^q \{2c^2 d f \lambda^{2q} \sin(\rho - \sigma) + bcd^2 \lambda^q \sin 2(\rho - \sigma) + bc f^2 \lambda^q \sin 2\rho + 2b^2 d f \sin(\rho + \sigma)\}].$$

Both ϵ_1 and J are dominated by one phase as far as $p \geq 2$ and $q \geq 1$. Therefore, the leptogenesis is linked strongly with the CP violation in the neutrino oscillations except for the case of $p = 1$ and $q = 0$.

In the case of the textures (A23) and (A24), ϵ_1 and J are given by putting $d = 0$ and $c = 0$ in eqs.(24) and (25), respectively. Therefore ϵ_1 and J are simply given as

$$\epsilon_1 \simeq \frac{3m_{D0}^2}{16\pi v^2} c^2 \lambda^4 \sin 2\rho, \quad (26)$$

$$J \simeq \frac{1}{64} a^2 b^4 c^2 f^4 \lambda^2 \frac{\Delta m_{\text{atm}}^2}{\Delta m_{\text{sol}}^2} \sin 2\rho,$$

for (A23), and

$$\epsilon_1 \simeq -\frac{3m_{D0}^2}{16\pi v^2} \frac{d^2 f^2}{b^2 + d^2} \lambda^4 \sin 2\sigma,$$




Figure 1: Predicted Y_B versus M_0 for $\sin 2\rho = 0.2, 0.5, 1$ in the texture A12. The gray region is allowed by the observed baryon asymmetry.

(27)

$$J \simeq \frac{1}{64} a^2 b^6 d^2 f^2 \lambda^2 \frac{\Delta m_{\text{atm}}^2}{\Delta m_{\text{sol}}^2} \sin 2\sigma ,$$

for (A24), respectively. The relative sign of ϵ_1 and J is same for (A23) while opposite for (A24) as seen in eqs.(26) and (27). Therefore, these textures are testable in the future.

In order to calculate the baryon asymmetry, we need the dilution factor involves the integration of the full set of Boltzmann equations [33]. A simple approximated solution which has been frequently used is given by [34]

$$\kappa = \frac{0.3}{K_R} (\ln K_R)^{-0.6} , \quad (10 \leq K_R \leq 10^6) , \quad (28)$$

where the parameter K_R is defined as the ratio of the thermal average of the N_1 decay rate and the Hubble parameter at the temperature $T = M_1$,

$$K_R = \frac{M_P}{1.7 \times 8\pi v^2 \sqrt{g_*}} \frac{(\overline{m}_D^\dagger \overline{m}_D)_{11}}{M_1} , \quad (29)$$

where $M_P \simeq 1.22 \times 10^{19}$ GeV is the Planck mass. The value of K_R is approximately ~ 23 , which is independent of M_1 , in our textures. Therefore, the dilution factor is not so suppressed.

By using this approximate dilution factor, we have calculated Y_B in our textures. Since $\epsilon_1 \simeq \lambda^4$ in all cases of our textures, we calculate Y_B for the case of (A12) with $f = 1$ in eq.(20)

as a typical one. Our numerical result is shown in figure 1. As seen in figure 1, the predicted Y_B is consistent with the observed value $Y_B = (1.7 \sim 8.1) \times 10^{-11}$ for $M_0 = 10^{13} \sim 10^{14} \text{GeV}$ and $\sin 2\rho = 0.2 \sim 1$. More detailed analyses will be given elsewhere. Thus, our textures of the Dirac neutrinos is favored in the scenario of the leptogenesis.

The discussion and summary are given as follows.

We have studied the leptogenesis in the neutrino mass matrix with two zeros, which reduces the number of CP violating phases. After see-saw realization in the textures A_1 and A_2 , we have found that one texture with three zeros has three phases, three textures with four zeros have two phases and two textures with five zeros have only one phase for the Dirac neutrino mass matrices, respectively. Taking a typical mass hierarchy of the right-handed Majorana neutrinos, we have calculated the relevant quantities for the leptogenesis and the low energy CP violation, ϵ_1 and J , respectively. The leptogenesis is linked strongly with the CP violation in the neutrino oscillations in our textures although there are a few exceptional cases. We have also given the prediction Y_B , which is consistent with the observed value. We will present more systematic and detailed analyses elsewhere. In order to test the link between the leptogenesis and the low energy CP violation, we expect that the measurement of the CP violation in the experiments of the long baseline neutrino oscillations.

We are grateful to T. Yanagida and P. H. Frampton for important comments on the number of phases in textures. We also thank T. Endoh, S. K. Kang and T. Morozumi for useful discussions at the YITP-W-02-05 on "Flavor mixing, CP violation and Origin of matter". This research is supported by the Grant-in-Aid for Science Research, Ministry of Education, Science and Culture, Japan (No.12047220).

References

- [1] Super-Kamiokande Collaboration, Y. Fukuda et al, Phys. Rev. Lett. **81** (1998) 1562; ibid. **82** (1999) 2644; ibid. **82** (1999) 5194.
- [2] Super-Kamiokande Collaboration, S. Fukuda et al. Phys. Rev. Lett. **86** (2001) 5651; 5656.
- [3] SNO Collaboration: Q. R. Ahmad et al., Phys. Rev. Lett. **87** (2001) 071301; nucl-ex/0204008, 0204009.
- [4] L. Wolfenstein, Phys. Rev. **D17** (1978) 2369;
S. P. Mikheyev and A. Yu. Smirnov, Yad. Fiz. **42** (1985) 1441.
- [5] G. L. Fogli, E. Lisi, D. Montanino and A. Palazzo, Phys. Rev. **D64** (2001) 093007;
J. N. Bahcall, M. C. Gonzalez-Garcia and C. Peña-Garay, JHEP **0108** (2001) 014;
M. Fukugita and M. Tanimoto, Phys. Lett. **B515** (2001) 30;
A. Bandyopadhyay, S. Choubey, S. Goswami and K. Kar, Phys. Lett. **B519** (2001) 83;
V. Barger, D. Marfatia and K. Whisnant, Phys. Rev. Lett. **88** (2002) 011302.
- [6] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870.
- [7] P.H. Frampton, S.L. Glashow and D. Marfatia, Phys. Lett. **B536** (2002) 79.
- [8] Z. Xing, Phys. Lett. **B530** (2002) 159; Phys. Lett. **B539** (2002) 85.
- [9] B. R. Desai, D. P. Roy and A. R. Vaucher, hep-ph/0209035.
- [10] P.H. Frampton, M.C. Oh and T. Yoshikawa, Phys. Rev. **D66** (2002) 033007.
- [11] A. Kageyama, S. Kaneko, N. Simoyama and M. Tanimoto, Phys. Lett. **B538** (2002) 96.
- [12] S.K. Kang and C.S. Kim, Phys. Rev. **D63** (2001) 113010.
- [13] Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, Proceedings of the Workshop, Stony Brook, New York, 1979, edited by P. van Nieuwenhuizen and D. Freedmann, North-Holland, Amsterdam, 1979, p.315;

T. Yanagida, in *Proceedings of the Workshop on the Unified Theories and Baryon Number in the Universe*, Tsukuba, Japan, 1979, edited by O. Sawada and A. Sugamoto, KEK Report No. 79-18, Tsukuba, 1979, p.95.

- [14] M. Fukugita and T. Yanagida. Phys. Lett. **B175** (1986) 45.
- [15] M. Flanz, E. A. Paschos and U. Sarkar, Phys. Lett. **B345** (1995) 248;
 W. Buchmüller and M. Plümacher, Phys. Lett. **B389** (1996) 73; *ibid.* **B431** (1998) 354;
 Int. J. Mod. Phys. **A15** (2000) 5047;
 A. Pilaftsis, Phys. Rev. **D56** (1997) 5431.
- [16] A. S. Joshipura, E. A. Paschos and W. Rodejohann, JHEP **0108** (2001) 029;
 W. Buchmüller and D. Wyler, Phys. Lett. **B 521** (2001) 291;
 J. Ellis and M. Raidal, hep-ph/0206174;
 S. Davidson and A. Ibarra, hep-ph/0206304;
 G. Branco, T. Morozumi, B. Nobre and M.N. Rebelo, Nucl. Phys. **B 617** (2001) 475;
 G. Branco, R. González Felipe, F. R. Joaquim and M. N. Rebelo, Nucl. Phys. **B 640** (2002) 202.
- [17] D. Falcone, hep-ph/0204335;
 Z. Xing, hep-ph/0206245;
 M. Fujii, K. Hamaguti and T. Yanagida, Phys. Rev. **D65** (2002) 115012;
 W. Rodejohann, Phys. Lett. **B542** (2002) 100.
- [18] M. Tanimoto, Phys. Rev. **D55** (1997) 322; Prog. Theor. Phys. **97** (1997) 901; Phys. Lett. **B435** (1998)373; Phys. Lett. **B462** (1999) 115;
 J. Arafune and J. Sato, Phys. Rev. **D55** (1997)1653;
 J. Arafune, M. Koike and J. Sato, Phys. Rev. **D56** (1997) 3093;
 H. Minakata and H. Nunokawa, Phys. Lett. **B413** (1997) 369; Phys. Rev. **D57** (1998) 4403.
 For recent proposal of the search for CP violation in neutrino oscillation, see Y. Itow et.al., hep-ex/0106019.
- [19] P. Frampton, S. Glashow and T. Yanagida, hep-ph/0208157.

- [20] T. Endoh, S. Kaneko, S. K. Kang, T. Morozumi and M. Tanimoto, hep-ph/0209020.
- [21] P. H. Chankowski and Z. Pluciennik, Phys. Lett. **B316** (1993) 312;
 K. S. Babu, C. N. Leung and J. Pantaleone, Phys. Lett. **B319** (1993) 191;
 M. Tanimoto, Phys. Lett. **360B** (1995) 41;
 P. H. Chankowski, W. Krolikowski and S. Pokorski, Phys. Lett. **B473** (2000) 109;
 J. Ellis, G. K. Leontaris, S. Lola and D. V. Nanopoulos, Eur. Phys. J. **C9** (1999) 389;
 J. Ellis and S. Lola, Phys. Lett. **B458** (1999) 310;
 J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. **B556** (1999) 3; JHEP **9909** (1999) 015; Nucl. Phys. **B569** (2000) 82;
 M. Carena, J. Ellis, S. Lola and C. E. M. Wagner, Eur. Phys. J. **C12** (2000) 507;
 N. Haba, Y. Matsui, N. Okamura and M. Sugiura, Eur. Phys. J. **C10** (1999) 677; Prog. Theor. Phys. **103** (2000) 145;
 N. Haba and N. Okamura, Eur. Phys. J. **C14** (2000) 347;
 N. Haba, N. Okamura and M. Sugiura, Prog. Theor. Phys. **103** (2000) 367;
 S. Antusch, M. Drees, J. Kersten, M. Lindner and M. Ratz, Phys. Lett. **B519** (2001) 238; **B525** (2002) 130;
 S. Antusch and M. Ratz, JHEP **0207** (2002) 059.
- [22] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57** (1986) 961.
- [23] J. Hisano, T. Moroi, K. Tobe, M. Yamaguchi and T. Yanagida, Phys. Lett. **B357** (1995) 579;
 J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. **D53** (1996) 2442.
- [24] J. Hisano, D. Nomura and T. Yanagida, Phys. Lett. **B437** (1998) 351;
 J. Hisano and D. Nomura, Phys. Rev. **D59** (1999) 116005;
 M.E. Gomez, G.K. Leontaris, S. Lola and J. D. Vergados, Phys. Rev. **D59** (1999) 116009;
 W. Buchmüller, D. Delepine and F. Vissani, Phys. Lett. **B459** (1999) 171;
 W. Buchmüller, D. Delepine and L.T. Handoko, Nucl. Phys. **B576** (2000) 445;
 J. Ellis, M.E. Gomez, G.K. Leontaris, S. Lola and D.V. Nanopoulos, Eur. Phys. J. **C14** (2000) 319;

- J. L. Feng, Y. Nir and Y. Shadmi, Phys. Rev. **D61** (2000) 113005;
S. Baek, T. Goto, Y. Okada and K. Okumura, Phys. Rev. **D63** (2001) 051701.
- [25] J. Sato, K. Tobe, and T. Yanagida, Phys. Lett. **B498** (2001) 189;
J. Sato and K. Tobe, Phys. Rev. **D63** (2001) 116010;
S. Lavignac, I. Masina and C. A. Savoy, Phys. Lett. **B520** (2001) 269.
- [26] J. A. Casas and A. Ibarra, Nucl. Phys. **B618** (2001) 171.
- [27] A. Kageyama, S. Kaneko, N. Simoyama and M. Tanimoto, Phys. Lett. **B527** (2002) 206;
Phys. Rev. **D65** (2002) 096010.
- [28] MEGA Collaboration, M. L. Brooks et al., Phys. Rev. Lett. **83** (1999) 1521.
- [29] J. Ellis, J. Hisano, M. Raidal and Y. Shimizu, hep-ph/0206110.
- [30] L. Covi, E. Roulet and F. Vissani, Phys. Lett. **B 384** (1996) 169.
- [31] J. A. Harvey and M. S. Turner, Phys. Rev. **D42** (1990) 3344;
S. Y. Khlebnikov and S. E. Shaposhnikov, Nucl. Phys. **B308** (1998) 169.
- [32] C. Jarlskog, Phys. Rev. Lett. **55** (1985) 1039.
- [33] M. Luty, Phys. Rev. **D45** (1992) 455;
M. Plümacher, Z. Phys. **C74** (1997) 549.
- [34] E. W. Kolb, M. S. Turner, *The early universe*, Redwood City, USA: Addison-Wesley (1990), (Frontiers in physics, 69);
M. Flanz and E. A. Paschos, Phys. Rev. **D58** (1998) 113009;
A. Pilaftsis, Int. J. Mod. Phys. **A14** (1999) 1811;
H. B. Nielsen and Y. Takanishi, Phys. Lett. **B 507** (2001) 241.